

## ORIGINAL ARTICLE

## A common variant near *TGFBR3* is associated with primary open angle glaucoma

Zheng Li<sup>1,3,†</sup>, R. Rand Allingham<sup>4,†</sup>, Masakazu Nakano<sup>5,†</sup>, Liyun Jia<sup>8,†</sup>, Yuhong Chen<sup>9,†</sup>, Yoko Ikeda<sup>6</sup>, Baskaran Mani<sup>1,11,13</sup>, Li-Jia Chen<sup>14</sup>, Changwon Kee<sup>15</sup>, David F. Garway-Heath<sup>16</sup>, Sarangapani Sripriya<sup>17</sup>, Nobuo Fuse<sup>18</sup>, Khaled K. Abu-Amero<sup>19,20</sup>, Chukai Huang<sup>21</sup>, Prasanthi Namburi<sup>22</sup>, Kathryn Burdon<sup>23,24</sup>, Shamira A. Perera<sup>1,13,2</sup>, Puya Gharahkhani<sup>25</sup>, Ying Lin<sup>26,27</sup>, Morio Ueno<sup>6</sup>, Mineo Ozaki<sup>28</sup>, Takanori Mizoguchi<sup>29</sup>, Subbiah Ramasamy Krishnadas<sup>30</sup>, Essam A. Osman<sup>19</sup>, Mei Chin Lee<sup>1</sup>, Anita S.Y. Chan<sup>1,2</sup>, Liza-Sharmini A. Tajudin<sup>31</sup>, Tan Do<sup>32</sup>, Aurelien Goncalves<sup>33</sup>, Pascal Reynier<sup>34</sup>, Hong Zhang<sup>35</sup>, Rupert Bourne<sup>36</sup>, David Goh<sup>2</sup>, David Broadway<sup>37</sup>, Rahat Husain<sup>2</sup>, Anil K. Negi<sup>38</sup>, Daniel H Su<sup>2</sup>, Ching-Lin Ho<sup>2</sup>, Augusto Azuara Blanco<sup>39</sup>, Christopher K.S. Leung<sup>14</sup>, Tina T. Wong<sup>1,13,2</sup>, Azhany Yakub<sup>31</sup>, Yutao Liu<sup>40,41</sup>, Monisha E. Nongpiur<sup>1,11,13</sup>, Jong Chul Han<sup>15</sup>, Do Nhu Hon<sup>32</sup>, Balekudaru Shantha<sup>42</sup>, Bowen Zhao<sup>8</sup>, Jinghong Sang<sup>8</sup>, NiHong Zhang<sup>8</sup>, Ryuichi Sato<sup>5</sup>, Kengo Yoshii<sup>7</sup>, Songhomita Panda-Jonas<sup>43</sup>, Allison E. Ashley Koch<sup>40</sup>, Leon W. Herndon<sup>4</sup>, Sayoko E. Moroi<sup>44</sup>, Pratap Challa<sup>4</sup>, Jia Nee Foo<sup>3</sup>, Jin-Xin Bei<sup>46,47</sup>, Yi-Xin Zeng<sup>46,47</sup>, Cameron P. Simmons<sup>48,49</sup>, Tran Nguyen Bich Chau<sup>48</sup>, Philomenadin Ferdinamarie Sharmila<sup>17</sup>, Merwyn Chew<sup>1</sup>, Blanche Lim<sup>1</sup>, Pansy O.S. Tam<sup>14</sup>, Elaine Chua<sup>1</sup>, Xiao Yu Ng<sup>1</sup>, Victor H.K. Yong<sup>1</sup>, Yaan Fun Chong<sup>1</sup>, Wee Yang Meah<sup>3</sup>, Saravanan Vijayan<sup>22</sup>, Sohn Seongsoo<sup>15</sup>, Wang Xu<sup>12</sup>, Yik Ying Teo<sup>3,12</sup>, Jessica N. Cooke Bailey<sup>50</sup>, Jae H. Kang<sup>51</sup>, Jonathan L. Haines<sup>50</sup>, Ching Yu Cheng<sup>1,11,13,2</sup>, Seang-Mei Saw<sup>1,12</sup>, E-Shyong Tai<sup>52</sup>, ICAARE-Glaucoma Consortium<sup>‡</sup>, NEIGHBORHOOD Consortium<sup>‡</sup>, Julia E. Richards<sup>44,45</sup>, Robert Ritch<sup>53</sup>, Douglas E. Gaasterland<sup>54</sup>,

<sup>†</sup>These authors contributed equally to this work.

<sup>‡</sup>A list of members of each consortium appears in the Supplementary Material, Note.

<sup>¶</sup>These authors also contributed equally to this work.

Received: December 1, 2014. Revised: March 9, 2015. Accepted: April 8, 2015

© The Author 2015. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

Louis R. Pasquale<sup>51,55</sup>, Jianjun Liu<sup>3</sup>, Jost B. Jonas<sup>43</sup>, Dan Milea<sup>1,13,2</sup>, Ronnie George<sup>42</sup>, Saleh A. Al-Obeidan<sup>19</sup>, Kazuhiko Mori<sup>6</sup>, Stuart Macgregor<sup>25</sup>, Alex W. Hewitt<sup>23,56</sup>, Christopher A. Girkin<sup>57</sup>, Mingzhi Zhang<sup>21</sup>, Periasamy Sundaresan<sup>22</sup>, Lingam Vijaya<sup>42</sup>, David A. Mackey<sup>23,58</sup>, Tien Yin Wong<sup>1,2,11</sup>, Jamie E. Craig<sup>24</sup>, Xinghuai Sun<sup>9,10,59</sup>, Shigeru Kinoshita<sup>6</sup>, Janey L. Wiggs<sup>57,¶</sup>, Chiea-Chuen Khor<sup>3,11,¶,\*</sup>, Zhenglin Yang<sup>26,27,60,¶</sup>, Chi Pui Pang<sup>14,¶</sup>, Ningli Wang<sup>8,¶</sup>, Michael A. Hauser<sup>4,40,¶</sup>, Kei Tashiro<sup>5,¶</sup>, Tin Aung<sup>1,2,11,13,¶</sup> and Eranga N. Vithana<sup>1,11,13,¶,\*</sup>

<sup>1</sup>Singapore Eye Research Institute, <sup>2</sup>Singapore National Eye Center, Singapore, Singapore, <sup>3</sup>Division of Human Genetics, Genome Institute of Singapore, Singapore, Singapore, <sup>4</sup>Department of Ophthalmology, Duke University Eye Center, Durham, NC, USA, <sup>5</sup>Department of Genomic Medical Sciences, <sup>6</sup>Department of Ophthalmology, <sup>7</sup>Department of Medical Statistics, Kyoto Prefectural University of Medicine, Kyoto, Japan, <sup>8</sup>Beijing Ophthalmology & Visual Sciences Key Laboratory, Beijing Tongren Eye Centre, Beijing Tongren Hospital, Capital Medical University, Beijing, China, <sup>9</sup>Department of Ophthalmology and Visual Science, Eye and ENT Hospital, Shanghai Medical School, <sup>10</sup>State Key Laboratory of Medical Neurobiology, Institutes of Brain Science, Fudan University, Shanghai, China, <sup>11</sup>Department of Ophthalmology, Yong Loo Lin School of Medicine, <sup>12</sup>Saw Swee Hock School of Public Health, National University of Singapore, Singapore, Singapore, <sup>13</sup>Duke-NUS Graduate Medical School, Singapore, Singapore, <sup>14</sup>Department of Ophthalmology and Visual Sciences, The Chinese University of Hong Kong Eye Hospital, Hong Kong, China, <sup>15</sup>Department of Ophthalmology, Samsung Medical Center, Sungkyunkwan University, School of Medicine, Seoul, Seoul Korea, <sup>16</sup>National Institute for Health Research Biomedical Research Centre at Moorfields Eye Hospital NHS Foundation Trust and University College London Institute of Ophthalmology, London, UK, <sup>17</sup>SNONGC Department of Genetics and Molecular Biology, Vision Research Foundation, Sankara Nethralaya, Chennai, India, <sup>18</sup>Department of Integrative Genomics, Tohoku Medical Megabank Organization, Sendai, Japan, <sup>19</sup>Department of Ophthalmology, College of Medicine, King Saud University, Riyadh, Saudi Arabia, <sup>20</sup>Department of Ophthalmology, College of Medicine, University of Florida, Jacksonville, FL, USA, <sup>21</sup>Chinese University of Hong Kong Joint Shantou International Eye Center, Shantou University, Shantou, China, <sup>22</sup>Department of Genetics, Aravind Medical Research Foundation, Madurai, Tamilnadu, India, <sup>23</sup>Menzies Institute for Medical Research, University of Tasmania, Hobart, Australia, <sup>24</sup>Department of Ophthalmology, Flinders University, Adelaide, Australia, <sup>25</sup>Department of Genetics and Computational Biology, Statistical Genetics, QIMR Berghofer Medical Research Institute, Brisbane, QLD, Australia, <sup>26</sup>Sichuan Provincial Key Laboratory for Human Disease Gene Study, Hospital of the University of Electronic Science and Technology of China and Sichuan Provincial People's Hospital, Chengdu, China, <sup>27</sup>School of Medicine, University of Electronic Science and Technology of China, Chengdu, China, <sup>28</sup>Ozaki Eye Hospital, 1-15, Kamezaki, Hyuga, Miyazaki 883-0066, Japan, <sup>29</sup>Mizoguchi Eye Hospital, 6-13 Tawara-machi, Sasebo, Nagasaki 857-0016, Japan, <sup>30</sup>Glaucoma Services, Aravind Eye Hospital, Madurai, Tamilnadu, India, <sup>31</sup>Department of Ophthalmology, School of Medical Sciences, Universiti Sains Malaysia, Kota Bharu, Kelantan, Malaysia, <sup>32</sup>Vietnam National Institute of Ophthalmology, Hanoi, Vietnam, <sup>33</sup>Ophthalmology Department, <sup>34</sup>Biochemistry Department, Angers University Hospital, Angers, France, <sup>35</sup>Department of Ophthalmology, Tongji Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China, <sup>36</sup>Huntingdon Glaucoma Diagnostic & Research Centre, Hinchingsbrooke Hospital, Huntingdon, UK, <sup>37</sup>Norfolk & Norwich University Hospital NHS Trust, Norwich, UK, <sup>38</sup>Heart of UK NHS Foundation Trust, Birmingham, UK, <sup>39</sup>School of Medicine, Dentistry and Biomedical Sciences, Centre for Experimental Medicine, Queen's University Belfast, Northern Ireland, UK, <sup>40</sup>Department of Medicine, Duke University Medical Center, Durham, NC, USA, <sup>41</sup>Department of Cellular Biology and Anatomy, Georgia Regents University, Augusta, Georgia, <sup>42</sup>Medical Research Foundation, Sankara Nethralaya, Chennai, India, <sup>43</sup>Department of Ophthalmology, Medical Faculty Mannheim of the Ruprecht-Karls-University Heidelberg, Heidelberg, Germany, <sup>44</sup>Department of Ophthalmology and Visual Sciences, <sup>45</sup>Department of Epidemiology, University of Michigan, Ann Arbor, MI, USA, <sup>46</sup>State Key Laboratory of Oncology in Southern China, Guangzhou, China, <sup>47</sup>Department of Experimental Research, Sun Yat-Sen

University Cancer Centre, Guangzhou, China, <sup>48</sup>Clinical Research Unit, Oxford University, 190 Ben Ham Tu, Ho Chi Minh City, Vietnam, <sup>49</sup>Centre for Tropical Medicine, Nuffield Department of Clinical Medicine, Oxford University, Oxford OX3 7LJ, UK, <sup>50</sup>Department of Epidemiology and Biostatistics, Case Western Reserve University, Cleveland, OH, USA, <sup>51</sup>Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA, USA, <sup>52</sup>Department of Medicine, National University Health System & National University of Singapore, Singapore, <sup>53</sup>Einhorn Clinical Research Center, Department of Ophthalmology, New York Eye and Ear Infirmary, New York, NY, USA, <sup>54</sup>Eye Doctors of Washington DC, Washington, DC, USA, <sup>55</sup>Department of Ophthalmology, Harvard Medical School, Massachusetts Eye and Ear Infirmary, Boston, MA, USA, <sup>56</sup>Centre for Eye Research Australia (CERA), University of Melbourne, Royal Victorian Eye and Ear Hospital, Melbourne, VIC, Australia, <sup>57</sup>Department of Ophthalmology, University of Alabama at Birmingham, Birmingham, AL, USA, <sup>58</sup>Centre for Ophthalmology and Visual Science, Lions Eye Institute, University of Western Australia, Perth, WA, Australia, <sup>59</sup>Myopia Key Laboratory of the Ministry of Health of China, Shanghai, China and <sup>60</sup>Chinese Academy of Sciences, Sichuan Translational Medicine Hospital, Chengdu, China

\*To whom correspondence should be addressed at: The Academia, 20 College Road, Discovery Tower level 6, Singapore 169856, Singapore. Tel: +65 65767216; Fax: +65 62252568. Email: eranga.n.v@sericom.sg (E. N. V.); 60 Biopolis Street, #02-01 Genome Building, Genome Institute of Singapore, Singapore 138672, Singapore. Tel: +65 68088200; Fax: +65 68088034. Email: khorcc@gis.a-star.edu.sg (C.-C. K.)

## Abstract

Primary open angle glaucoma (POAG), a major cause of blindness worldwide, is a complex disease with a significant genetic contribution. We performed Exome Array (Illumina) analysis on 3504 POAG cases and 9746 controls with replication of the most significant findings in 9173 POAG cases and 26 780 controls across 18 collections of Asian, African and European descent. Apart from confirming strong evidence of association at *CDKN2B-AS1* (rs2157719 [G], odds ratio [OR] = 0.71,  $P = 2.81 \times 10^{-33}$ ), we observed one SNP showing significant association to POAG (*CDC7-TGFBR3* rs1192415,  $OR_{G-allele} = 1.13$ ,  $P_{meta} = 1.60 \times 10^{-8}$ ). This particular SNP has previously been shown to be strongly associated with optic disc area and vertical cup-to-disc ratio, which are regarded as glaucoma-related quantitative traits. Our study now extends this by directly implicating it in POAG disease pathogenesis.

## Introduction

Glaucoma is the leading cause of irreversible visual impairment and blindness, affecting >60 million people worldwide, and it is estimated that the number of affected individuals will reach 80 million in 2020 (1–3). Primary open angle glaucoma (POAG) is the most prevalent form of glaucoma in most populations and is characterized by progressive retinal ganglion cell (RGC) loss that causes characteristic structural changes of the optic nerve with associated with visual field loss in the face of an open drainage angle in the eye. POAG has a strong genetic component that has been well documented (4). Indeed, several susceptibility loci have been identified for POAG through the use of linkage and association studies (5). Genes known to contribute to glaucoma include myocilin (*MYOC*), optineurin (*OPTN*), TANK-binding kinase 1 (*TBK1*) and WD repeat domain 36 (*WDR36*) (6–12). However, mutations in these genes account for no >5–10% of all POAG cases in the general population (5).

It is likely that POAG, as a complex trait, results from the interactions of multiple genes and environmental factors (13–15). Genome-wide association studies (GWAS) have provided further insights into the genetic basis of POAG (16). The first GWAS on POAG was conducted on 1263 POAG cases and 34 877 controls from Iceland. Genome-wide significant association was detected at the *CAV1-CAV2* locus on Chromosome 7q31 and subsequently was replicated in a multi-ethnic sample collection from Sweden, the UK, Australia, Hong Kong and China (17). This was rapidly followed by five other GWAS studies, which utilized either advanced or non-advanced POAG cases derived from populations

of European or East Asian ancestries (18–22). These latter studies led to the discovery of nine additional genetic regions associated with POAG disease risk (*TMCO1*, *CDKN2B-AS1*, *SIX1-SIX6*, an intergenic region on chromosome 8q22, *ABCA1*, *GAS7*, *AFAP1*, *GMDS* and *PMM2*). Several of these genetic loci have been replicated in ethnically diverse populations, demonstrating them to be bona fide POAG associations with global implications (23–25).

Similar in concept and laboratory chemistry to the whole-genome genotyping chip design, the exome array approach evaluates putative functional coding variants selected from the exome sequences of >12 000 individuals (26). In addition, the exome array also contains >5000 common variant SNPs from GWAS arrays with a minor allele frequency (MAF) exceeding 5% which can serve as ancestry informative markers. Exome array genotyping allows us to specifically explore the possible contribution of potentially functional coding variants in POAG disease susceptibility.

## Results

### Common genetic variants in *CDKN2B-AS1* and *TGFBR3-CDC7* are associated with POAG

We conducted a two-stage Exome Chip discovery and replication on POAG cases and normal controls. For the discovery stage (Stage 1), genotyping was performed using the Illumina Infinium HumanExomeBeadChip (v1.0) on a total of 3822 POAG cases and 10 426 normal controls drawn from seven countries (Table 1). In addition to the  $\approx 247\,000$  SNP markers present on the standard

**Table 1.** Sample collections of POAG cases and controls for Stages 1 (discovery) and 2 (replication)

Collection	N cases	N controls	Ethnicity	Age of cases	Age of controls <sup>a</sup>	Collection comment
Singapore	850	2347	Singaporean Chinese	71.6 ± 10.1	58.88 ± 9.6	New recruitment
Japan	923	640	Japanese	65.3 ± 13.1	71.9 ± 5.8	Previously GWAS in Nakano <i>et al.</i> (25)
USA-African-Americans	590	636	African American	65.4 ± 12.3	54.8 ± 9.8	New recruitment
China-Beijing	587	461	Northern Chinese	58.5 ± 12.5	Population-based controls	New recruitment
Hong Kong	375	2962	Southern Chinese	62.3 ± 15.3	Population-based controls	Previously described for replication in Thorleifsson <i>et al.</i> (17)
South India	121	716	Indian	60.9 ± 12.0	51.0 ± 6.5	New recruitment
Vietnam	58	1984	Vietnamese	63 ± 7.1	Population-based controls	New recruitment
Total discovery	3504	9746				2206 cases are new discovery samples
Stage 2 replication						
Singapore-2	520	5473	Singaporean Chinese	71.1 ± 10	Population-based controls	New recruitment
Japan-2	935	996	Japanese	64.3 ± 14.0	57.5 ± 13.9	N = 411 used for replication in Nakano <i>et al.</i> (25)
USA-African-American 2	497	304	African-American	69.1 ± 11.0	66.7 ± 13.1	New recruitment
South India-2	453	2496	Indian	62.5 ± 9.9	58.9 ± 10.1	New recruitment
Korea	400	454	Korean	59.0 ± 11.8	40.3 ± 14.1	New recruitment
Saudi Arabia	236	655	Middle Eastern	60.8 ± 12.7	54.4 ± 11.7	New recruitment
Malaysia	132	2540	Malay	65.1 ± 8.2	58.7 ± 11.0	New recruitment
China-Beijing 2	115	251	Northern Chinese	54.2 ± 12.4	71.53 ± 7.16	New recruitment
UK	336	6090	European	71.4 ± 10.8	Population-based controls	New recruitment
China-Shantou	247	289	Southern Chinese	52.9 ± 19.4	75.7 ± 6.1	Previously described for replication in Thorleifsson <i>et al.</i> (17)
Germany	56	142	European	67.9 ± 11.4	78.4 ± 8.9	New recruitment
Vietnam-2	76	245	Vietnamese	52.4 ± 17.4	51.3 ± 17.8	New recruitment
France	80	75	European	75.6 ± 8.5	73.5 ± 8.3	New recruitment
China-Shanghai, Chengdu 2	181	286	Southern Chinese	54.7 ± 16.5	84.7 ± 11.7	Previously described in Chen <i>et al.</i> (19)
China-Shanghai, Chengdu	608	1005	Southern Chinese	49.6 ± 17.0	62.9 ± 12.1	Previously described in Chen <i>et al.</i> (19)
USA (NEIGHBOR)	2170	2347	European descendant	66.4	68	Previously described in Wiggs <i>et al.</i> (22)
USA (GLAUGEN)	976	1140	European descendant	63.6	65.5	Previously described in Wiggs <i>et al.</i> (22)
Australia (ANZRAG)	1155	1992	European descendant	60.5 ± 14.3	55.6 ± 14.4	Previously described in Gharahkhani <i>et al.</i> (20)
Total replication	9173	26 780				3425 cases are new replication samples
Total all samples	12 677	36 526				5631 cases are new in this report.

<sup>a</sup>Population-based controls are ascertained from large-scale studies and do not have demographic data available. Based on many well-described examples, both by others and us, the frequency of POAG in the general population is uncommon (i.e. <5%). In this regard, the false-negative rate for POAG status in the population-based controls is likely to be low and thus the effect of loss of statistical power is negligible.

Illumina Exome array (26), we also included an extra 25 000 coding frame SNP markers obtained from exome sequencing of 2000 individuals of East Asian descent. Stringent quality control (QC) filters were applied to both SNPs and samples: per SNP call rate ≥99%, per-sample call rate ≥95%, non-monomorphic SNPs and non-significant deviation from Hardy-Weinberg equilibrium (HWE)  $P \geq 10^{-6}$ . Samples under suspicion of cross-contamination and biologically related samples were removed by verification of extreme heterozygosity and identical by descent/identical by state information, if applicable. We further performed principal component analysis (PCA) to verify that cases and controls were well-matched ancestrally (Supplementary Material, Fig. S1) (27–32). As a result, a total of 3504 POAG cases and 9746

controls passing QC filters were included for association analysis using unconditional logistic regression, with adjustments for the principal components (PCs). From these datasets, a total of 2206 POAG cases represent entirely new patient collections, which have not been previously reported (Supplementary Material, Table S1). Each study-specific point estimate was then summarized using fixed-effects meta-analysis ( $N_{\text{meta-analysis}} = 7$  collections). A quantile-quantile plot derived from the meta-analysis P-values showed no significant dispersion of test statistics from the expected distribution ( $\lambda_{\text{GC}} = 1.042$ ; Supplementary Material, Fig. S2) suggesting that the association results were not confounded by cryptic population stratification. Using additive effect models, we observed experiment-wide significant

( $P = 0.05/272\,000$  SNPs =  $1.84 \times 10^{-7}$ ) evidence of association at CDKN2B-AS1 (rs2157719 [G], per-allele OR = 0.73,  $P = 1.10 \times 10^{-7}$ ) (Supplementary Material, Fig. S3).


All SNP markers showing  $P < 0.0005$  in the discovery stage were followed up in a validation stage (Stage 2) comprised of up to 9173 POAG cases and 26 780 controls. A total of 21 SNPs at 20 independent loci were brought forward for validation genotyping using Sequenom MassARRAY iPLEX or *in silico* look-ups if genome-wide genotyping data were available (see Supplementary Material, Table S2).

CDKN2B-AS1 rs2157719 once again showed significant evidence of association in the meta-analysis of all replication collections (OR = 0.70,  $P = 2.48 \times 10^{-27}$ ) as well as meta-analysis of all samples tested (rs2157719 [G], OR = 0.71,  $P_{\text{meta}} = 2.81 \times 10^{-33}$ ) (Supplementary Material, Table S2 and Fig. S4). No heterogeneity of effect between Asians (OR = 0.72,  $P = 1.15 \times 10^{-15}$ ) and Europeans (OR = 0.69,  $P = 5.54 \times 10^{-19}$ ) were detected for this marker (Fig. 1A), consistent with the multiple previous reports describing association at this locus (18,22,25). Apart from CDKN2B-AS1 rs2157719, a second SNP marker (CDC7-TGFBR3 rs1192415) showed clear evidence of replication in Stage 2 (OR = 1.12,  $P = 1.04 \times 10^{-5}$ , Fig. 1B and Supplementary Material, Table S2). On meta-analysis with the exome-chip discovery findings, genome-wide significant evidence of association was observed for rs1192415 (OR = 1.13,  $P_{\text{meta}} = 1.6 \times 10^{-8}$ ) (Fig. 1B; Supplementary Material, Table S2). Although the risk allele ranged from between 11 and 34% across all ethnic groups studied for this marker, the association observed appeared to be uniform and consistent across most groups with little overall heterogeneity ( $I^2$  index

< 20%, Supplementary Material, Table S3). For rs1192514, the association appeared stronger in Asians (OR = 1.17,  $P = 1.48 \times 10^{-7}$ ) compared with Europeans (OR = 1.10,  $P = 0.01$ ) (Fig. 1B), although the difference was not statistically significant ( $P_{\text{het}} = 0.17$ ). One other marker (FNDC3B rs4894796) was nominally significant in the replication stage (OR = 0.95,  $P = 0.02$ ) and remained suggestively associated with POAG in the overall meta-analysis (OR = 0.93,  $P = 1.40 \times 10^{-5}$ ) (Fig. 1C; Supplementary Material, Table S2). However, the association appeared to be nearly entirely driven by the Asian POAG collections (OR = 0.89) compared with the Europeans (OR = 0.99,  $P_{\text{het}}$  between Asians and Europeans = 0.0042) (Fig. 1C). A recently reported GWAS on POAG showed strong association with common SNP markers mapping to AFAP1 (rs4478172) (20). Looking up on our exome dataset, we successfully genotyped rs7437940 ( $r^2 = 0.18$ ,  $D' = 0.97$  with rs4478172) which also mapped within AFAP1. We note significant association at this AFAP1 marker (rs7437940: Stage 1  $P = 1.94 \times 10^{-5}$ , Stage 2  $P = 0.08$ ,  $P$ -value for meta-analysis =  $4.25 \times 10^{-6}$ ) (Supplementary Material, Table S2), which supports the previous report (20).

### Expression of POAG-associated genes in ocular tissues

We examined the mRNA expression of CDC7, TGFBR3 and FNDC3B in multiple eye tissues. Expression of all three genes was observed in tissues relevant to POAG such as the trabecular meshwork, optic disc and nerve. In contrast to TGFBR3 and FNDC3B, which were expressed in all tested ocular tissues, CDC7 expression was absent in the iris, ciliary body and choroid (Fig. 2).



**Figure 1.** Forest plots showing evidence of association between SNPs: (A) CDKN2B-AS1 rs2157719, (B) CDC7/TGFBR3 rs1192415 and (C) FNDC3B rs4894796. The vertical line represents a per-allele odds ratio of 1.00. The oblongs represent point estimates (referring to the per-allele odds ratio), with the height of the oblongs inversely proportional to the standard error of the point estimates. Horizontal lines indicate the 95% confidence interval for each point estimate. Meta-analysis of Stages 1 and 2, OR,  $P_{\text{meta}}$  and  $I^2$  was labeled on the right-hand side for corresponding analysis. For rs4894796 genotyping, see Supplementary Material, Information for sample collections.





Figure 1 Continued

We also investigated the localization of FNDC3B protein in normal ocular tissues, focusing our attention on cornea and the tissues of the outflow pathway. FNDC3B could be immunolocalized to cells

in the trabecular meshwork and all three layers of the cornea (Fig. 2). Immunolocalization of TGFBR3 and CDC7 could not be similarly investigated due to lack of availability of specific antibodies.



**Figure 2.** Analysis of *FNDC3B*, *TGFBR3* and *CDC7* expression in ocular tissues. (A) The *FNDC3B*-specific 162 bp and *TGFBR3*-specific 152 bp amplification product was observed in all analyzed ocular tissues. *CDC7*-specific 242 bp product was observed in sclera, cornea, trabecular meshwork, retina, optic disc and optic nerve. The ubiquitously expressed gene, *ACTB* was used as the normalizing control. A no template sample acted as the negative control (NC) to ensure non-contamination of the RT-PCR reaction mix. The variable M denotes molecular-weight marker. (B) Immunoblot of whole cell lysates from NPCE, retinal pigment epithelial (ARPE19) and HTM cells, probed for *FNDC3B* and  $\beta$ -actin, as a loading control. Positions of the ~133 and ~44 kDa forms of *FNDC3B* are indicated. All ocular cells analyzed expressed the ~133 kDa protein, while ARPE19 cells expressed a smaller ~44 kDa isoform of *FNDC3B*. (C) Immunolocalization of *FNDC3B* in human eye tissues. Strong immunofluorescence labeling of *FNDC3B* (green) was seen in the ciliary muscle (CM) (top row). Scale bar: 100  $\mu$ m. In the trabecular meshwork (TM, middle row), *FNDC3B* (green) labeling was relatively weaker. Scale bar: 40  $\mu$ m. *FNDC3B* positive immunoreactivity was also observed in cornea epithelial (\*) and cornea endothelial cells (white arrows) (bottom row). Nuclei were stained with DAPI (blue). Scale bar: 100  $\mu$ m.

### Analysis of rare variants from the exome-chip discovery collection

We next proceeded to conduct gene-based tests on mutational load to further investigate the role of low-frequency variants in POAG for all patient collections in the discovery stage. Gene-based tests are an alternative to single-marker tests for association, which are often underpowered to detect association with rare variants. To more directly address the impact of low frequency, non-synonymous genetic variants, we

considered only the variants with MAF of <5% (33). As a result, we were able to assess a total of 7822 genes having at least two such variants using the sequence kernel association optimal test (34). We note two genes (*MYO18A* and *SULF2*) which showed nominal evidence of association on burden test ( $P = 6.26 \times 10^{-7}$  and  $P = 5.78 \times 10^{-5}$ , respectively), but these findings are primarily driven by the small Vietnamese collection ( $N = 58$  cases and  $N = 1984$  controls). Removal of this dataset resulted in a marked reduction of the burden test association results (Supplementary Material, Table S4).

## Discussion

The present study supports the association of POAG a locus that has been previously implicated with vertical cup-disc ratio, a glaucoma-associated endophenotype. This locus is defined by rs1192415, a common SNP marker mapping to the intergenic region between *CDC7* and *TGFBR3* (allele [G], OR = 1.13,  $P_{\text{meta}} = 1.6 \times 10^{-8}$ ). We were also able to strongly corroborate the established association of POAG with *CDKN2B-AS1.CDC7-TGFBR3* rs1192415 has previously been reported to be strongly associated with overall optic disc area, which together with VCDR is a glaucoma-related quantitative trait (35,36). Examination of this locus in >4700 POAG cases and >90 000 controls of European descent in a recent study showed direction of effect consistent with our data (OR = 1.06,  $P = 0.12$ ) (37). Indeed, in the European section of our study ( $N = 4773$  cases and 11786 controls), we note a similar effect size (OR = 1.10,  $P = 0.01$ ) in keeping with that shown by Springelkamp and others. The effect of this locus appeared to be much stronger in Asians (OR = 1.17,  $P = 1.48 \times 10^{-7}$ ), leading to a genome-wide significant association upon meta-analysis of all collections. Each copy of the rs1192415 minor allele is associated with a relatively modest increase in POAG risk of 1.13 at exome-wide significance. The consistency of the effect across 21 of 24 POAG collections ( $P_{\text{het}} = 0.22$ ,  $I^2$  index for heterogeneity = 16.9%), lend further credence to the observed association.

It is of note that the G allele of the rs1192415 is associated with increase in disc area and therefore a larger VCDR. This may indicate a lower threshold for diagnosing glaucoma in those individuals harboring the risk allele (38). However, as the diagnosis of POAG in this study was not based solely upon the presence of an increased cup-disc ratio but also compatible visual field loss, it is unlikely that selection bias has influenced the result obtained with rs1192415. We note that *FNDC3B* was one of 16 loci associated with central corneal thickness (sentinel SNP being rs4894535) in a meta-analysis conducted on >20 000 individuals of European and Asian descent (39). That study also showed marker rs4894535 to be associated with POAG in 2979 cases and 7399 controls of European descent (OR = 0.83,  $P = 5.6 \times 10^{-4}$ ). We were able to confirm association of rs4894535 with POAG in 5810 cases and 13 175 controls with readily available DNA for genotyping across 13 collections in this study (OR = 0.91,  $P = 0.0018$ ; Supplementary Material, Table S5), thus lending further support for this association.

It is noteworthy that the *FNDC3B* SNP we report here (rs4894796) was nominally associated with POAG in the overall meta-analyses (OR = 0.93,  $P = 1.4 \times 10^{-5}$ ). In particular, the association at rs4894796 was observed to be particularly strong in Asians (OR = 0.89,  $P = 7.93 \times 10^{-8}$ ) compared with Europeans (OR = 0.99,  $P = 0.71$ ). Unsurprisingly, rs4894796 was found within a different linkage disequilibrium (LD) block than the SNP associated with central corneal thickness and POAG (rs4894535). The low LD between rs4894796 and rs4894535 suggests that the association signal at the latter SNP is not likely to be driven by rs4894796. An in-depth inspection of the LD patterns of this genetic region between European and East Asian populations, which currently appear visually identical according to available heat maps (<http://www.hapmap.org>), may be useful to better understand the degree of LD dissimilarity between populations and its bearing on the results obtained for *FNDC3B*. Of note, a further SNP (rs6445055) at *FNDC3B* showed genome-wide significant association for intraocular pressure (IOP) ( $P = 4.9 \times 10^{-8}$ ), but only marginal association was seen for POAG per se ( $P = 0.03$ , OR = 0.92) in 4284 POAG cases and 95 560 controls of European descent (21). As the *FNDC3B* SNP associations with POAG are still suggestive rather than

affirmative it is presumptive at this stage to speculate that it could be one of the links for IOP-dependent mechanisms of POAG.

This is one of the largest studies on the genetics of POAG, yet the power to detect genes with small effects was limited. One crucial reason for this apparent loss of power could be the scope of the genetic content which was used for interrogation (as exomic content only comprises <2% of the entire human genome, much useful data could be missed if the bulk of the true positive genetic associations for POAG lie in the non-coding regions of the genome). The fact that our discovery cohort was not mono-ethnic may also have reduced the power to detect ethnic specific variants of small effect sizes due to significant differences in allele frequencies between ethnic groups. The phenotypic heterogeneity and the lack of standardized clinical criteria across the cohorts may also have contributed to a loss in power. The level of IOP is commonly used to subdivide POAG into two subtypes: POAG with high IOP (>21 mmHg; named high-tension glaucoma, HTG) and NTG with normal IOP (<21 mmHg). In our cohorts, we had differing number of these subtypes with some cohorts having more NTG or HTG than others. In this study, we also did not subdivide POAG subjects into HTG and NTG to discover subtype-specific variants, focusing instead on identifying genetic variants for the overarching phenotype of POAG. However, when *CDC7-TGFBR3* rs1192415 and *FNDC3B* rs4894796 were analyzed within NTG and HTG subgroups, they did not show differences in strength of association by subtype (data not shown).

All three loci we report in this study (*CDKN2B-AS1*, *TGFBR3-CDC7* and *FNDC3B*) contain genes which may contribute to the regulation of transforming growth factor- $\beta$  (TGF- $\beta$ ) signaling. TGF- $\beta$  has been implicated previously in glaucomatous optic nerve damage and RGC death (40–42). The TGF- $\beta$  family includes TGF $\beta$ 1, TGF $\beta$ 2 and TGF $\beta$ 3, all of which bind to TGF- $\beta$  receptor type-2 (TGFBR2). All TGF- $\beta$  family members are dimeric polypeptide growth factors that inhibit the progression of cell cycle, which in turn may lead to terminal differentiation or apoptosis (40,43,44). TGF- $\beta$  also modulates developmental and repair processes in several tissues. TGF- $\beta$  signaling has been implicated in a wide variety of diseases including inflammation, autoimmune disorders, fibrosis, cancer, cataracts as well as glaucoma (40,43,44). The most strongly POAG-associated locus, *CDKN2B-AS1*, has been shown to regulate the transcription of cyclin-dependent kinase inhibitor 2A and 2B (*CDKN2A* and *CDKN2B*) (45), which inhibit cell proliferation via the TGF- $\beta$  pathway by inducing G1-phase cell cycle arrest (44). Burdon *et al.* reported the up-regulation of *CDKN2A* and *CDKN2B* in response to elevated IOP (18). Collectively, these data suggest a link between the most well-recognized physiological risk factor of POAG and a downstream molecular response that may lead to RGC death. In this context, both *CDC7* and *TGFBR3* are also of interest and could have relevance to glaucomatous optic nerve damage and RGC death.

*CDC7* encodes a cell division cycle protein with kinase activity that also interacts with *CDKN2A*. *TGFBR3* is a TGF- $\beta$  super family co-receptor, and is the most abundant of all TGF- $\beta$  receptors (46). Through protein crystallography, murine TGF- $\beta$  3 ZP domain (ZP-C) has been recently identified as a novel major TGF- $\beta$ -binding site (47). It has been suggested that *TGFBR3* may serve to enhance the binding of TGF- $\beta$  ligands to TGF- $\beta$  type II receptors by binding TGF- $\beta$  and presenting it to TGFBR2, the receptor for all three TGF- $\beta$  ligands. A linkage between *FNDC3B*, an oncogene and TGF- $\beta$  signaling was also reported recently by Cai *et al.* Overexpression of *FNDC3B* was shown to induce epithelial-to-mesenchymal transition and activate several cancer pathways, including PI3-kinase/



Akt, Rb1 and TGF- $\beta$  signaling (48). FNDC3B also induced expression of all three TGF- $\beta$  ligands and promoted TGFBR1 cell-surface localization (48). This connection of POAG-associated genes/loci with the TGF- $\beta$  signaling pathway therefore lends further credence to the hypothesis of TGF $\beta$  pathway involvement in glaucoma.

However, despite these attractive speculations on the genes vicinal to the associated SNPs, it is possible that these variants may also affect distant as yet unidentified target genes. Definitive evidence for the involvement of these genes in the pathogenesis of POAG awaits confirmation in other datasets, as well as the identification and characterization of functional variants.

## Materials and Methods

### Sample collections

#### Ethics statement

Ethics approval was obtained from the Centralized Institutional Review Board for the Singaporean patient sample and data collection and for the conduct of this study. All study protocols for patient and control sample collection were approved by the respective relevant Medical Ethics Committees of each participating site. All studies were conducted under the tenets of the Declaration of Helsinki with written informed consent obtained from all participants.

#### POAG and health controls subjects inclusion criteria

POAG cases in this study were defined by the following criteria: the presence of glaucomatous optic neuropathy (defined as loss of neuroretinal rim with a vertical cup : disc ratio of  $>0.7$  or an inter-eye asymmetry of  $>0.2$  and/or notching attributable to glaucoma) with compatible visual field loss, open angles on gonioscopy, and absence of secondary causes of glaucomatous optic neuropathy. POAG patients with a mean IOP without treatment that is consistently  $<21$  mmHg on diurnal testing are classified as having NTG, whereas those with a mean IOP without treatment that is consistently  $>21$  mmHg are classified as having HTG. Patients who were unable to give informed consent, or with secondary glaucoma due to trauma, uveitis, neovascularization, pseudoexfoliation, pigment dispersion, etc., were excluded from this study.

Controls in this study were recruited in a hospital-based or population-based manner. Hospital-based controls were all generally over the age of 40 years and confirmed to have no sign of glaucoma or other major eye diseases except for mild cataract and mild refractive errors (defined as  $|SE| < 3D$ ) by an ophthalmic examination. These subjects at time of recruitment had IOP of  $<21$  mmHg with open angles, healthy optic nerves, normal visual fields, and no family history of glaucoma. Population-based controls were ethnically matched healthy individuals over the age of 40 years, unless indicated otherwise (see Table 1; Supplementary Material for more details of the samples used in this study).

### Genotyping and data QC

Study participants in the discovery stage were genotyped using the Illumina's Infinium HumanExomeBeadChip (Version 1.0) + Semi-Custom BeadChip (Illumina Inc.) that contains ~250 000 SNPs of base content and an additional 25 000 East Asian-specific polymorphisms located on the coding frame. Stringent QC filters were applied after the laboratory work in genotyping completed. SNP markers that had missingness exceeding 5%, gross departure from HWE ( $P$ -value  $< 1e-6$ ) or were monomorphic were excluded from subsequent analysis. Likewise, individual samples with an

overall call rate  $<95\%$  were excluded. Samples were subjected to biological relationship verification by using the principle of variability in allele sharing according to the degree of relationship. Identity-by-state information was derived by PLINK. Those individuals who showed evidence of cryptic relatedness were removed before PC analysis was conducted. In addition, samples showing gender discrepancies between the clinical gender and genetically inferred gender were removed. A total of 1008 samples were excluded after rigorous application of QC filters (568 sample exclusions were due to PCA ancestral outliers, 246 were excluded due to per-sample call rate  $<95\%$ , 168 were excluded due to first-degree familial relationships detected from the discovery stage exome-chip genotyping, and 26 were excluded due to suspicions of sample contamination). PC analysis was undertaken using EIGENSTRAT to account for spurious associations resulting from ancestral differences of individual SNPs (49). PCs showing significant effect on univariate analysis were used to correct for any underlying population substructure. We adjusted for the top three PCs (PC1–PC3) for Singapore, Hong Kong, Japan, USA-African Americans, China-Beijing and Vietnam. The Indian POAG collection was adjusted for the top 10 PCs (PC1–PC10), as there was more population substructure in this collection. After adjustment, we observed minimal evidence of genomic inflation ( $\lambda_{GC} = 1.042$ ), thereby suggesting that this well-described method of controlling for population stratification was adequate in our study. Genotyping clouds for the key SNPs CDKN2B-AS1 rs2157719, CDC7-TGFB3rs1192415 and FNDC3Brs4894796 were directly visualized (Supplementary Material, Fig. S5) to ensure good quality.

For Stage 2 (replication stage), genotyping was performed using the Sequenom MassArray platform (www.sequenom.com). Samples recruited at the latter stage of replication were genotyped for the key SNPs CDC7-TGFB3rs1192415, and FNDC3Brs4894796 using pre-developed Taqman Assays (Applied Biosystems, Foster City, CA, USA; www.appliedbiosystems.com)

### Statistical analysis

We contrasted the genotypes between POAG cases and healthy controls via single-SNP analysis using unconditional logistic regression fitted for genotype trend effects (1-degree-of-freedom score test). To do this, the PLINK software [version 1.07] (50) was used for modeling within a logistic regression framework, adjusting for age, gender and genetic ancestry (reflected by PCs). Manhattan and LD plots were created using Haploview [version 3.2] (51). Q-Q and regional association plots were created using the software R [www.r-project.org] (52). Meta-analysis summarizing the results across all cohorts was performed using both fixed and random-effects modeling weighted in an inverse-variance manner (53). This method weighs each study according to effective sample size and cohort-specific MAF of the associated variants. To avoid an otherwise unacceptable number of false positive signals as an artifact of multiple testing, the threshold for exome-wide significance,  $P < 2 \times 10^{-7}$ , was considered to be statistically significant. Heterogeneity of the meta-analyses was calculated by measuring  $I^2$ .

### Expression analysis of genes

Expression of CDC7, TGFB3 and FNDC3B was assessed by semi-quantitative reverse transcription-PCR (RT-PCR) using gene-specific primers (CDC7-forward 5'-TTTCTCCCCAGCGTGACC-3', CDC7-reverse 5'-GCAATTTCTCTCAGGTCCTAC-3'; TGFB3-forward 5'-TCTCCTCAGTCCACATCCAC-3', TGFB3-reverse 5'-TGC

TGATGAAAACCTGGACCAC-3'; FNDC3B-forward 5'-AGCATCATCT TCCCACACA-3', FNDC3B-reverse 5'-AAGAAGGAGGGCTGTTG AGG-3') on total RNA extracted from a variety of ocular tissues (cornea, sclera, retina and retinal pigment epithelium, iris, lens capsule and optic nerve) as described earlier (54). We used the ubiquitously expressed *ACTB* gene (forward 5'-CCAACCGCGA GAAGATGA-3' and reverse 5'-CCAGAGCGGTACAGGGATAG-3') as amplification and normalizing control.

### Western blotting

Cell lines obtained from American Type Culture Collection (Manassas, VA, USA) were the human retinal pigment epithelial cell line (APRE19), human Trabecular Meshwork cell line (HTM) was purchased from ScienCell Research Laboratories (Carlsbad, CA, USA) and the human non-pigmented ciliary epithelial cell line (NPCE) is a kind gift from Prof. Miguel Coca-Prados from Yale School of Medicine. Cell lysates were obtained by lysing individual cell lines with lysis buffer (50 mM Tris-HCl, pH 8, 150 mM NaCl, 1.0% Nonidet P-40, 0.5% deoxycholate, 0.1% SDS, 0.2 mM NaVO<sub>4</sub>, 10 mM NaF, 0.4 mM EDTA and 10% glycerol). SDS-PAGE resolved proteins were transferred to Hybond-C Extra nitrocellulose membranes (Amersham Life Science Inc., Arlington Heights, IL, USA). Membranes were blocked and blotted by 5% nonfat milk, 0.1% Tween 20 in Tris-buffered saline (20 mM Tris-HCl, pH 7.6, 150 mM NaCl) for 1 h before incubation with FNDC3B for 1 h (1:250) (Sigma-Aldrich Corp., St. Louis, MO, USA). Actin-horseradish peroxidase (HRP) (1:50 000) from Santa Cruz Biotechnology (Dallas, TX, USA). The bound primary antibodies were detected by horseradish peroxidase-conjugated secondary antibodies (GE Healthcare Biosciences, Pittsburgh, PA, USA), and visualized by Luminata Forte Western HRP substrate (Millipore, Bedford, MA, USA).

### Immunofluorescence confocal microscopy

Immunofluorescence confocal microscopy was performed on antigen retrieved 4 µm paraffin sections. Blocking of tissue sections was performed with blocking buffer (5% nonfat milk, 5% FBS, 0.1% PBS-Tween; 1× pen/strep) for 1 h at RT. FNDC3B antibody (Sigma-Aldrich Corp.) was diluted (1:50) into blocking buffer and incubated overnight at 4°C. Secondary FITC (1:300)-labelled anti-rabbit antibody (Jackson Laboratories, Westgrove, PA, USA) was also diluted in blocking buffer and incubated at RT for 1 h followed by application of Vectashield with 4',6-diamidino-2-phenyl-indole (DAPI) (Vector Laboratories, Burlingame, CA, USA). Coverslips were then used to overlay the sections and stored in the dark at 4°C until viewing with Olympus Fluoview 1000 confocal microscope (Olympus Optical Co. Ltd., Tokyo, Japan).

### Supplementary Material

Supplementary Material is available at HMG online.

### Acknowledgements

We thank all the study participants, the staff from all involved studies and sites who contributed to this study. The authors J.E.C., K.P.B., D.A.M., and others acknowledge the support of Ms Bronwyn Usher-Ridge in patient recruitment and data collection, and Dr Patrick Danoy and Dr Johanna Hadler for genotyping of the ANZRAG cohort. They also acknowledge David C. Whiteman and Graham Radford-Smith for providing access to the control samples used with the ANZRAG glaucoma cases. The authors

D.M. thank Dr Philippe Gohier and Dr Ghislaine Jallet for patient recruitment and data collection and acknowledge the technical assistance provided by the Centre de Recherche Biologique (Dr Odile Blanchet) and Centre de Recherche Clinique (Pr Marc-Antoine Custeau), University Hospital, Angers, France. The Singapore study of E.N.V., T.A. and C.C.K was supported by a Biomedical Research Council (BMRC) grant in Singapore, Ref: BMRC 10/1/35/19/675. This research was also partly supported by a grant (NMRC/TCR/008-SERI/2013) from the Singapore National Research Foundation under its Translational and Clinical Research Flagship Programme and administered by the Singapore Ministry of Health's National Medical Research Council. We also acknowledge the following source of funding support for recruitment and genotyping of population-based cohorts SIMES and SCES: National Medical Research Council, Singapore (NMRC/TCR/002-SERI/2008, (R626/47/2008TCR), CSA R613/34/2008, NMRC 0796/2003, STaR/0003/2008), the National Research Foundation of Singapore, the Biomedical Research Council, Singapore (BMRC 09/1/35/ 19/616 and 08/1/35/19/550) and Genome Institute of Singapore (GIS/12-AR2105). The Singapore Tissue Network and the Genome Institute of Singapore, Agency for Science, Technology and Research, Singapore provided services. The research of M.A.H. was supported by NIH R01 EY023646, EY13315 (MAH) and by P30-EY005722. The Japanese sample collection from Kyoto was supported by the grants from the Collaborative Development of Innovative Seeds of Japan Science and Technology Agency (JST) to M.K. and K.T., from the Ministry of Health, Labor and Welfare of Japan to M.N., K.M., K.T. and S.K., and from Santen Pharmaceutical Co. Ltd. to S.K. and K.T. The collection of Japanese POAG cases by N.F. was supported by a grant provided by the Ministry of Health, Labor and Welfare of Japan. The glaucoma research of WNL is supported by National Natural Science Foundation of China project (81030016). The research of Y. Chen was supported by National Natural Science Foundation of China (81200723) and glaucoma research of Xinghuai Sun was supported by Special Scientific Research Project of Health Professions (201302015). This research project was supported by the National Natural Science Foundation of China (81170883 (Z.Y.) and 81430008 (Z.Y.)). The glaucoma research of C.P.P. is supported by the Health and Medical Research Fund (HMRP, Ref: 01122236 and 11120801), Hong Kong and the General Research Fund from the Research Grants Council (grant number 468810), Hong Kong. Support for recruitment of Australian & New Zealand Registry of Advanced Glaucoma (ANZRAG) was provided by the Royal Australian and New Zealand College of Ophthalmology (RANZCO) Eye Foundation. Genotyping was funded by the National Health and Medical Research Council of Australia (#535074 and #1023911). This work was also supported by funding from NHMRC #1031362 awarded to Jamie E. Craig, NHMRC #1037838 awarded to Alex W. Hewitt, NHMRC #1048037 awarded to Stuart L. Graham, NHMRC #1009844 awarded to Robert J. Casson and Ivan Goldberg, NHMRC #1031920 and an Alcon Research Institute grant awarded to David A. Mackey, an Allergan Unrestricted grant awarded to Andrew J. White, the BrightFocus Foundation and a Ramaciotti Establishment Grant. S.M. is supported by Australian Research Council (ARC) and Australian National Health & Medical Research Council (NHMRC) Fellowships. Controls for the ANZRAG discovery cohort were drawn from the Australian Cancer Study, the Study of Digestive Health, and from a study of inflammatory bowel diseases. The Australian Cancer Study was supported by the Queensland Cancer Fund and the National Health and Medical Research Council (NHMRC) of Australia (Program no. 199600, awarded to David C. Whiteman, Adele C. Green, Nicholas K. Hayward, Peter

G. Parsons, David M. Purdie and Penelope M. Webb, and program number 552429, awarded to David C. Whiteman). The Study of Digestive Health was supported by grant number 5 RO1 CA 001833 from the National Cancer Institute (awarded to David C. Whiteman). The Barrett's and Esophageal Adenocarcinoma Genetic Susceptibility Study (BEAGESS) sponsored the genotyping of oesophageal cancer and Barrett's oesophagus cases, which were used as unscreened controls in the ANZRAG discovery cohort. BEAGESS was funded by grant R01 CA136725 from the National Cancer Institute Collection of Saudi POAG cases and controls was supported by the Glaucoma Research Chair at College of Medicine, King Saud University, Riyadh, Saudi Arabia. NEI Glaucoma Human Genetics Collaboration (NEIGHBOR): genotyping services for the NEIGHBOR study were provided by the CIDR and were supported by the NEI through grant HG005259-01 (J.L.W.). Additionally, CIDR is funded through a federal contract from the NIH to The Johns Hopkins University, contract number HHSN268200782096C. Collecting and processing samples for the NEIGHBOR data set was supported by the NEI through American Recovery and Reinvestment Act (ARRA) grants 3R01EY015872-05S1 (J.L.W.) and 3R01EY019126-02S1 (M.A.H.). Genotype imputation and meta-analysis were supported by EY022305 (J.L.W.). Funding for the collection of cases and controls was provided by the following NIH grants: EY015543 (R.R. Allingham); EY006827 (D. Gaasterland); HL73042, HL073389, EY13315, EY023646 (M.A.H.); CA87969, CA49449, UM1 CA167552 (J.H. Kang); EY009149 (P.R. Lichter); HG004608 (C. McCarty); EY008208 (F.A. Medeiros); EY015473 (L.R.P.); EY012118 (M. Pericak-Vance); EY015682 (A. Realini); EY011671, EY09580 (J.E. Richards); EY013178 (J.S. Schuman); RR015574, EY015872, EY010886, EY009847, EY014104 (J.L.W.); EY011008, EY144428, EY144448 and EY18660 (K. Zhang). J.L.W. and L.R.P. are also supported by the Harvard Glaucoma Center for Excellence and Research to Prevent Blindness. J.N.C.B. is supported by NIH T32 EY007157 (CWRU) and T32 EY21453-2 (VUMC). MEEI case-control sample: genotyping for the Massachusetts Eye and Ear Infirmary (MEEI) case-control sample was performed at the Broad Institute of MIT and Harvard with funding support from the NIH GEI (Gene Environment Initiative) (U01HG04424 and U01HG004728). The GENEVA Coordinating Center (U01HG004446) assisted with genotype cleaning. Imputation was supported by NIH EY022305 (JLW). Collection of cases and controls was supported by NIH EY015872 (JLW) and NIH P30 014104 (JLW). Funding to pay the Open Access publication charges for this article was provided by Singapore Eye Research Institute.

*Conflict of Interest statement.* None declared.

## References

- Quigley, H.A. (1996) Number of people with glaucoma worldwide. *Br. J. Ophthalmol.*, **80**, 389–393.
- Quigley, H.A. and Broman, A.T. (2006) The number of people with glaucoma worldwide in 2010 and 2020. *Br. J. Ophthalmol.*, **90**, 262–267.
- Tham, Y.C., Li, X., Wong, T.Y., Quigley, H.A., Aung, T. and Cheng, C.Y. (2014) Global prevalence of glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-analysis. *Ophthalmology*, **121**, 2081–2090.
- Allingham, R.R., Liu, Y. and Rhee, D.J. (2009) The genetics of primary open-angle glaucoma: a review. *Exp. Eye Res.*, **88**, 837–844.
- Fan, B.J., Wang, D.Y., Lam, D.S. and Pang, C.P. (2006) Gene mapping for primary open angle glaucoma. *Clin. Biochem.*, **39**, 249–258.
- Sheffield, V.C., Stone, E.M., Alward, W.L., Drack, A.V., Johnson, A.T., Streb, L.M. and Nichols, B.E. (1993) Genetic linkage of familial open angle glaucoma to chromosome 1q21-q31. *Nat. Genet.*, **4**, 47–50.
- Stone, E.M., Fingert, J.H., Alward, W.L., Nguyen, T.D., Polansky, J.R., Sundén, S.L., Nishimura, D., Clark, A.F., Nystuen, A., Nichols, B.E. et al. (1997) Identification of a gene that causes primary open angle glaucoma. *Science (New York, NY)*, **275**, 668–670.
- Rezaie, T., Child, A., Hitchings, R., Brice, G., Miller, L., Coca-Prados, M., Heon, E., Krupin, T., Ritch, R., Kreutzer, D. et al. (2002) Adult-onset primary open-angle glaucoma caused by mutations in optineurin. *Science (New York, NY)*, **295**, 1077–1079.
- Sarfaraizi, M., Child, A., Stoilova, D., Brice, G., Desai, T., Trifan, O.C., Poinosawmy, D. and Crick, R.P. (1998) Localization of the fourth locus (GLC1E) for adult-onset primary open-angle glaucoma to the 10p15-p14 region. *Am. J. Hum. Genet.*, **62**, 641–652.
- Monemi, S., Spaeth, G., DaSilva, A., Popinchalk, S., Ilitchev, E., Liebmann, J., Ritch, R., Heon, E., Crick, R.P., Child, A. et al. (2005) Identification of a novel adult-onset primary open-angle glaucoma (POAG) gene on 5q22.1. *Hum. Mol. Genet.*, **14**, 725–733.
- Fingert, J.H. (2011) Primary open-angle glaucoma genes. *Eye (London, UK)*, **25**, 587–595.
- Fingert, J.H., Robin, A.L., Stone, J.L., Roos, B.R., Davis, L.K., Scheetz, T.E., Bennett, S.R., Wassink, T.H., Kwon, Y.H., Alward, W.L. et al. (2011) Copy number variations on chromosome 12q14 in patients with normal tension glaucoma. *Hum. Mol. Genet.*, **20**, 2482–2494.
- Fan, B.J., Leung, Y.F., Wang, N., Lam, S.C., Liu, Y., Tam, O.S. and Pang, C.P. (2004) Genetic and environmental risk factors for primary open-angle glaucoma. *Chin. Med. J.*, **117**, 706–710.
- Wiggs, J.L., Damji, K.F., Haines, J.L., Pericak-Vance, M.A. and Allingham, R.R. (1996) The distinction between juvenile and adult-onset primary open-angle glaucoma. *Am. J. Hum. Genet.*, **58**, 243–244.
- Kang, J.H., Wiggs, J.L., Rosner, B.A., Hankinson, S.E., Abdrou, W., Fan, B.J., Haines, J. and Pasquale, L.R. (2010) Endothelial nitric oxide synthase gene variants and primary open-angle glaucoma: interactions with sex and postmenopausal hormone use. *Invest. Ophthalmol. Visual Sci.*, **51**, 971–979.
- Hirschhorn, J.N. and Daly, M.J. (2005) Genome-wide association studies for common diseases and complex traits. *Nat. Rev.*, **6**, 95–108.
- Thorleifsson, G., Walters, G.B., Hewitt, A.W., Masson, G., Helgason, A., DeWan, A., Sigurdsson, A., Jonasdottir, A., Gudjonsson, S.A., Magnusson, K.P. et al. (2010) Common variants near CAV1 and CAV2 are associated with primary open-angle glaucoma. *Nat. Genet.*, **42**, 906–909.
- Burdon, K.P., Macgregor, S., Hewitt, A.W., Sharma, S., Childlow, G., Mills, R.A., Danoy, P., Casson, R., Viswanathan, A.C., Liu, J.Z. et al. (2011) Genome-wide association study identifies susceptibility loci for open angle glaucoma at TMCO1 and CDKN2B-AS1. *Nat. Genet.*, **43**, 574–578.
- Chen, Y., Lin, Y., Vithana, E.N., Jia, L., Zuo, X., Wong, T.Y., Chen, L.J., Zhu, X., Tam, P.O., Gong, B. et al. (2014) Common variants near ABCA1 and in PMM2 are associated with primary open-angle glaucoma. *Nat. Genet.*, **46**, 1115–1119.
- Gharahkhani, P., Burdon, K.P., Fogarty, R., Sharma, S., Hewitt, A.W., Martin, S., Law, M.H., Cremin, K., Bailey, J.N., Loomis, S.J. et al. (2014) Common variants near ABCA1, AFAP1 and GMD5

- confer risk of primary open-angle glaucoma. *Nat. Genet.*, **46**, 1120–1125.
21. Hysi, P.G., Cheng, C.Y., Springelkamp, H., Macgregor, S., Bailey, J.N., Wojciechowski, R., Vitart, V., Nag, A., Hewitt, A.W., Hohn, R. et al. (2014) Genome-wide analysis of multi-ancestry cohorts identifies new loci influencing intraocular pressure and susceptibility to glaucoma. *Nat. Genet.*, **46**, 1126–1130.
  22. Wiggs, J.L., Yaspan, B.L., Hauser, M.A., Kang, J.H., Allingham, R.R., Olson, L.M., Abdrabou, W., Fan, B.J., Wang, D.Y., Brodeur, W. et al. (2012) Common variants at 9p21 and 8q22 are associated with increased susceptibility to optic nerve degeneration in glaucoma. *PLoS Genet.*, **8**, e1002654.
  23. Fan, B.J., Wang, D.Y., Pasquale, L.R., Haines, J.L. and Wiggs, J.L. (2011) Genetic variants associated with optic nerve vertical cup-to-disc ratio are risk factors for primary open angle glaucoma in a US Caucasian population. *Invest. Ophthalmol. Visual Sci.*, **52**, 1788–1792.
  24. Cao, D., Jiao, X., Liu, X., Hennis, A., Leske, M.C., Nemesure, B. and Hejtmancik, J.F. (2012) CDKN2B polymorphism is associated with primary open-angle glaucoma (POAG) in the Afro-Caribbean population of Barbados, West Indies. *PLoS ONE*, **7**, e39278.
  25. Nakano, M., Ikeda, Y., Tokuda, Y., Fuwa, M., Omi, N., Ueno, M., Imai, K., Adachi, H., Kageyama, M., Mori, K. et al. (2012) Common variants in CDKN2B-AS1 associated with optic-nerve vulnerability of glaucoma identified by genome-wide association studies in Japanese. *PLoS ONE*, **7**, e33389.
  26. Huyghe, J.R., Jackson, A.U., Fogarty, M.P., Buchkovich, M.L., Stancakova, A., Stringham, H.M., Sim, X., Yang, L., Fuchsberger, C., Cederberg, H. et al. (2013) Exome array analysis identifies new loci and low-frequency variants influencing insulin processing and secretion. *Nat. Genet.*, **45**, 197–201.
  27. (ANZgene), Australia and New Zealand Multiple Sclerosis Genetics Consortium. (2009) Genome-wide association study identifies new multiple sclerosis susceptibility loci on chromosomes 12 and 20. *Nat. Genet.*, **41**, 824–828.
  28. Kooner, J.S., Saleheen, D., Sim, X., Sehmi, J., Zhang, W., Frossard, P., Been, L.F., Chia, K.S., Dimas, A.S., Hassanali, N. et al. (2011) Genome-wide association study in individuals of South Asian ancestry identifies six new type 2 diabetes susceptibility loci. *Nat. Genet.*, **43**, 984–989.
  29. Reveille, J.D., Sims, A.M., Danoy, P., Evans, D.M., Leo, P., Pointon, J.J., Jin, R., Zhou, X., Bradbury, L.A., Appleton, L.H. et al. (2010) Genome-wide association study of ankylosing spondylitis identifies non-MHC susceptibility loci. *Nat. Genet.*, **42**, 123–127.
  30. Hoglinger, G.U., Melhem, N.M., Dickson, D.W., Sleiman, P.M., Wang, L.S., Klei, L., Rademakers, R., de Silva, R., Litvan, I., Riley, D.E. et al. (2011) Identification of common variants influencing risk of the tauopathy progressive supranuclear palsy. *Nat. Genet.*, **43**, 699–705.
  31. Khor, C.C., Davila, S., Breunis, W.B., Lee, Y.C., Shimizu, C., Wright, V.J., Yeung, R.S., Tan, D.E., Sim, K.S., Wang, J.J. et al. (2011) Genome-wide association study identifies FCGR2A as a susceptibility locus for Kawasaki disease. *Nat. Genet.*, **43**, 1241–1246.
  32. Sawcer, S., Hellenthal, G., Pirinen, M., Spencer, C.C., Patsopoulos, N.A., Moutsianas, L., Dilthey, A., Su, Z., Freeman, C., Hunt, S.E. et al. (2011) Genetic risk and a primary role for cell-mediated immune mechanisms in multiple sclerosis. *Nature*, **476**, 214–219.
  33. Bonnefond, A., Skrobek, B., Lobbens, S., Eury, E., Thuillier, D., Cauchi, S., Lantieri, O., Balkau, B., Riboli, E., Marre, M. et al. (2013) Association between large detectable clonal mosaicism and type 2 diabetes with vascular complications. *Nat. Genet.*, **45**, 1040–1043.
  34. Lee, S., Wu, M.C. and Lin, X. (2012) Optimal tests for rare variant effects in sequencing association studies. *Biostatistics*, **13**, 762–775.
  35. Ramdas, W.D., van Koolwijk, L.M., Ikram, M.K., Jansonius, N. M., de Jong, P.T., Bergen, A.A., Isaacs, A., Amin, N., Aulchenko, Y.S., Wolfs, R.C. et al. (2010) A genome-wide association study of optic disc parameters. *PLoS Genet.*, **6**, e1000978.
  36. Khor, C.C., Ramdas, W.D., Vithana, E.N., Cornes, B.K., Sim, X., Tay, W.T., Saw, S.M., Zheng, Y., Lavanya, R., Wu, R. et al. (2011) Genome-wide association studies in Asians confirm the involvement of ATOH7 and TGFBR3, and further identify CARD10 as a novel locus influencing optic disc area. *Hum. Mol. Genet.*, **20**, 1864–1872.
  37. Springelkamp, H., Höhn, R., Mishra, A., Hysi, P.G., Khor, C.C., Loomis, S.J., Bailey, J.N., Gibson, J., Thorleifsson, G., Janssen, S.F. et al. (2015) Meta-analysis of genome-wide association studies identifies novel loci that influence cupping and the glaucomatous process. *Nat. Commun.*, **5**, 4883.
  38. Garway-Heath, D.F., Ruben, S.T., Viswanathan, A. and Hitchings, R.A. (1998) Vertical cup/disc ratio in relation to optic disc size: its value in the assessment of the glaucoma suspect. *Br. J. Ophthalmol.*, **82**, 1118–1124.
  39. Lu, Y., Vitart, V., Burdon, K.P., Khor, C.C., Bykhovskaya, Y., Mirshahi, A., Hewitt, A.W., Koehn, D., Hysi, P.G., Ramdas, W. D. et al. (2013) Genome-wide association analyses identify multiple loci associated with central corneal thickness and keratoconus. *Nat. Genet.*, **45**, 155–163.
  40. Fuchshofer, R. (2011) The pathogenic role of transforming growth factor-beta2 in glaucomatous damage to the optic nerve head. *Exp. Eye Res.*, **93**, 165–169.
  41. Pasquale, L.R., Dorman-Pease, M.E., Luttly, G.A., Quigley, H.A. and Jampel, H.D. (1993) Immunolocalization of TGF-beta 1, TGF-beta 2, and TGF-beta 3 in the anterior segment of the human eye. *Invest. Ophthalmol. Visual Sci.*, **34**, 23–30.
  42. Pena, J.D., Taylor, A.W., Ricard, C.S., Vidal, I. and Hernandez, M.R. (1999) Transforming growth factor beta isoforms in human optic nerve heads. *Br. J. Ophthalmol.*, **83**, 209–218.
  43. Massague, J., Blain, S.W. and Lo, R.S. (2000) TGFbeta signaling in growth control, cancer, and heritable disorders. *Cell*, **103**, 295–309.
  44. Hannon, G.J. and Beach, D. (1994) p15INK4B is a potential effector of TGF-beta-induced cell cycle arrest. *Nature*, **371**, 257–261.
  45. Pasmant, E., Sabbagh, A., Vidaud, M. and Bieche, I. (2011) ANRIL, a long, noncoding RNA, is an unexpected major hotspot in GWAS. *FASEB J.*, **25**, 444–448.
  46. Blobel, G.C., Liu, X., Fang, S.J., How, T. and Lodish, H.F. (2001) A novel mechanism for regulating transforming growth factor beta (TGF-beta) signaling. Functional modulation of type III TGF-beta receptor expression through interaction with the PDZ domain protein, GIPC. *J. Biol. Chem.*, **276**, 39608–39617.
  47. Diestel, U., Resch, M., Meinhardt, K., Weiler, S., Hellmann, T. V., Mueller, T.D., Nickel, J., Eichler, J. and Muller, Y.A. (2013) Identification of a novel TGF-beta-binding site in the Zona Pellucida C-terminal (ZP-C) domain of TGF-beta-receptor-3 (TGFR-3). *PLoS ONE*, **8**, e67214.
  48. Cai, C., Rajaram, M., Zhou, X., Liu, Q., Marchica, J., Li, J. and Powers, R.S. (2012) Activation of multiple cancer pathways and tumor maintenance function of the 3q amplified oncogene FNDC3B. *Cell Cycle (Georgetown, Tex)*, **11**, 1773–1781.
  49. Price, A.L., Patterson, N.J., Plenge, R.M., Weinblatt, M.E., Shadick, N.A. and Reich, D. (2006) Principal components analysis

- corrects for stratification in genome-wide association studies. *Nat. Genet.*, **38**, 904–909.
50. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller, J., Sklar, P., de Bakker, P.I., Daly, M.J. *et al.* (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.*, **81**, 559–575.
51. Barrett, J.C., Fry, B., Maller, J. and Daly, M.J. (2005) Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics (Oxford, England)*, **21**, 263–265.
52. Ihaka, R. and Gentleman, R. (1996) R: a language for data analysis and graphics. *J. Comput. Graph. Stat.*, **5**, 299–314.
53. Nalls, M.A., Plagnol, V., Hernandez, D.G., Sharma, M., Sheerin, U.M., Saad, M., Simon-Sanchez, J., Schulte, C., Lesage, S., Sveinbjornsdottir, S. *et al.* (2011) Imputation of sequence variants for identification of genetic risks for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet*, **377**, 641–649.
54. Vithana, E.N., Aung, T., Khor, C.C., Cornes, B.K., Tay, W.T., Sim, X., Lavanya, R., Wu, R., Zheng, Y., Hibberd, M.L. *et al.* (2011) Collagen-related genes influence the glaucoma risk factor, central corneal thickness. *Hum. Mol. Genet.*, **20**, 649–658.